

STI/STD (Sexually Transmitted Infection/ Disease)	Signs and Symptoms	How it Spread?	How can you prevent it?
<p>Chlamydia (bacterial)</p>	<ul style="list-style-type: none"> • Most people don't show signs/symptoms. • Symptoms may show up 7-21 days after having sex. • Pain or burning when you urinate (pee). • Discharge from the vagina, penis, or rectum (butt) that smells or looks strange. • Bleeding from the vagina during/after sex. • Pain or swelling in the testicles. • Pain or bleeding from the rectum. 	<ul style="list-style-type: none"> • By having vaginal, anal, or oral sex without a condom with someone who has chlamydia. • It can pass to a baby during birth. 	<ul style="list-style-type: none"> • Use condoms or dental dams. • Limit your sex partners. • Get tested in all the places (vagina, penis, anus, and throat) you have sex. • Don't have sex until you and your partner(s) finish treatment. • If pregnant, seek early prenatal care to prevent health problems for you and your baby.
	<ul style="list-style-type: none"> • Most people don't show signs/symptoms. • Symptoms may show up right away or as 		<ul style="list-style-type: none"> • Use condoms or dental dams.

Gonorrhea (bacterial)

long as 30 days after having sex.

- Pain or burning when you urinate (pee).

- Fluid from the vagina, penis, or rectum

(butt) that smells or looks strange.

- Itching, soreness, or bleeding in the rectum

or painful bowel movements (poop).

- Fever

- Bleeding from the vagina during/after sex.

- Pain or swelling in the testicles.

- Soreness or redness in the

- By having vaginal, anal, or oral sex

without a condom with someone who

has gonorrhea.

- It can pass to a baby during birth.

- Limit your sex partners.

- Get tested in all the places

(vagina, penis, anus, and

throat) you have sex.

- Don't have sex until you and

your partner(s) finish treatment.

- If pregnant, seek early prenatal

care to prevent health problems

for you and your baby.

Syphilis

- Most people don't show signs/symptoms.

- Symptoms may show up within 10-90 days.

- Symptoms can include chancre (sore where

the bacteria entered – vagina, anus, lips, or

- By having vaginal, anal, or oral sex

without a condom or sometimes

- Use condoms or dental dams.

- Limit your sex partners.

- Get tested regularly.

- Don't have sex until you and

(bacterial)

mouth; rash on one or more areas of the

body (back, hands and feet); hair loss; wart like bumps; smooth, white patches inside

the mouth or tongue.

- Symptoms can disappear and come back.

genital touching with someone with

syphilis.

- It can pass to a baby during pregnancy or birth.

your partner(s) finish treatment.

- If pregnant, seek early prenatal

care to prevent health problems

for you and your baby.